


## Deep Vein Thrombosis Assessment and Referral Form

Incorporating NICE Clinical Guideline 144 (2012)

First Name.....	Gender M/F	Date .....
Last Name.....	Telephone Number.....	
Address.....		
Hospital Number: RXR.....	NHS Number: .....	
GP.....	Religion.....	Completed by: .....

TWO-LEVEL DVT WELLS SCORE		
Clinical feature	Points	Score
Active cancer (treatment ongoing, within 6/12 or palliative)	1	
Paralysis, paresis or recent plaster immobilisation of the lower extremities	1	
Recently bedridden for 3 days or more or major surgery within 12 weeks requiring general or regional anaesthesia	1	
Localised tenderness along the distribution of the deep venous system	1	
Entire leg swollen	1	
Calf swelling at least 3 cm larger than asymptomatic side	1	
Pitting oedema confined to the symptomatic leg	1	
Collateral superficial veins (non-varicose)	1	
Previously documented DVT	1	
An alternative diagnosis is at least as likely as DVT (See Alternate Diagnoses Table below)	-2	
DVT Clinical probability simplified score		
DVT likely	2 points or more	
DVT unlikely	1 point or less	

Calf Measurements	
Left	cm
Right	cm
Blood Results	
D-Dimer	
Plts	
INR	
WCC	
Patient's Weight	
Kg	
Tinzaparin 20,000 IU/ml (Px on ED Card)	
Dose	mls
Time	
Weekend Tinzaparin	
Weekend doses prescribed on ED Card for patient to return. <input type="checkbox"/>	
OR: Take-home prescription provided for self-administration. <input type="checkbox"/>	
OR: Take-home prescription provided and DN contacted. <input type="checkbox"/>	



Alternative Diagnoses	Features	Tick/Cross
Superficial thrombo-phlebitis <b>&gt;Still need to rule out DVT</b>	Painful, tender, subcutaneous cord-like vein	
Bruising	Hx of injury, colour change, localised tenderness	
Haematoma	Local trauma, swelling, tenderness, fluctuation, hx of anticoagulation	
Muscle tear	Hx injury, specific muscle tenderness	
Ruptured Baker's Cyst	Previous swelling, arthritis, diffuse tenderness	
Cellulitis	Redness, fever, lymphangitis	
Compartment syndrome	Hx injury, severe pain, colour changes, diminished pulses	

\*In the 'DVT Likely' patient the D-Dimer is required only for decision-making in the DVT Clinic and for their audit purposes.

\*\*Refer to ELHT Tinzaparin Prescribing Advice Poster for Contraindications and Dosing. (In clean utility rooms.)