



Think Ectopic



- **ECTOPIC PREGNANCY - 12,000 cases** a year in the UK
- Vaginal bleeding, shoulder tip pain, lateralising pain, syncope are **RED FLAGS** for imminent or actual rupture
- Do a pregnancy test in all women of reproductive age with suspected symptoms attributable to an ectopic – some women are **unaware** they are pregnant
- Women with suspected ectopic require **urgent** Gynaecology assessment
- In women with pain/bleeding in early pregnancy where the risk of an ectopic is considered low, offer an early EPAU appointment & **written safety netting advice.**

HSIB investigation found communication and advice provided about Early Pregnancy Clinics could be improved. Link to the report:

(<https://www.hssib.org.uk/patient-safety-investigations/the-diagnosis-of-ectopic-pregnancy/investigation-report/>)

The RCEM has a publicly available discharge advice leaflet template, with thanks to Dr Hannah Bell, via this link: ([https://res.cloudinary.com/studio-republic/images/v1670604364/Patient Discharge Advice Colour/Patient Discharge Advice Colour.pdf? i=AA](https://res.cloudinary.com/studio-republic/images/v1670604364/Patient%20Discharge%20Advice%20Colour/Patient%20Discharge%20Advice%20Colour.pdf?i=AA))

For Safety Alerts and RCEM issued Safety Flashes see:

www.rcem.ac.uk/Safety